



Carotid Endarterectomy Instructions

1. Carotid Endarterectomy

- Keep drain site covered and dry for 48 hours.
- Make sure you turn your neck as much as possible.
- If your wound becomes swollen or bulging call the office immediately.
- If you experience any weakness on the side opposite of the operation, call the office immediately.
- If you experience any loss of vision on the side of the operation, call the office immediately.
- Monitor your blood pressure at home if you have a blood pressure cuff. If your blood pressure is below 100/60, hold your blood pressure medicine and call your Cardiologist or Internist.
-

1. Dressing Care

- **Dermabond® Dressing Care (superglue without a dressing):**
 - OK to shower day of surgery.
 - Lift nothing heavier than gallon milk jug or 5lbs for 1 week.

OR

- **General Wound Care Instructions for Closed Incisions with dressings:**
 - Dressing should be kept dry for 48 hours.
 - Remove outer dressing and OK to shower in 48 hours.
 - If you have staples, they will be removed in 1-2 weeks after the operation in the office.
 - Leave Steri-Strips in place.
 - No soaking in the tub or swimming for 1 week.

Call The Crawford Clinic right away if:

- You develop a fever over 101.5°F (38.5°C).
- You have redness, swelling, pain, or drainage from the incisions.
- You become dizzy and faint.
- You have chest pain.
- You become short of breath.
- You have abdominal pain or swelling that gets worse.

Call during office hours if:

- You have questions about the procedure or its result.
- You want to make another appointment.
- If you require more pain medication.